

(19) 日本国特許庁 (JP)

(12) 公開特許公報(A)

(11) 特許出願公開番号

特開2004-171

(P2004-171A)

(43) 公開日 平成16年1月8日 (2004.1.8)

(51) Int. Cl. ⁷

A23L 1/30

F1

A23L 1/30

B

テーマコード (参考)

4B018

審査請求 未請求 請求項の数 5 O L (全 14 頁)

(21) 出願番号 特願2003-81157 (P2003-81157)
 (22) 出願日 平成15年3月24日 (2003.3.24)
 (31) 優先権主張番号 特願2002-82036 (P2002-82036)
 (32) 優先日 平成14年3月22日 (2002.3.22)
 (33) 優先権主張国 日本国 (JP)

特許法第30条第1項適用申請有り 平成13年10月
 1日 トワ商事株式会社頒分の「マカ・エキスパウダー」
 に発表

(71) 出願人 300022526
 トワ商事株式会社
 東京都新宿区上落合2-11-1 橋本ビル
 (74) 代理人 100062764
 弁理士 樺澤 襄
 (74) 代理人 100092565
 弁理士 樺澤 聡
 (74) 代理人 100112449
 弁理士 山田 哲也
 (72) 発明者 小川 博
 大阪府富田林市藤沢台1-2-102
 (72) 発明者 光永 俊郎
 京都府京都市左京区下鴨膳部町101-3

最終頁に続く


(54) 【発明の名称】 マカを含有した機能性食品

(57) 【要約】

【課題】 シワや白髪の増加、体力および生殖機能の低下に有効と考えられる血中成長ホルモン量上昇作用および持久力増加作用を有する機能性食品を提供する。

【解決手段】 マカの根の乾燥物の6.0%以上8.0%以下の含水アルコール抽出物を含有する。血中成長ホルモン量上昇作用および持久力増加作用を有する。成長ホルモンの分泌が減少し、血中成長ホルモン量が低下したようなときに、5週間以上継続して摂取する。老化現象の抑制を期待できる。

【選択図】 図1



【特許請求の範囲】

【請求項 1】

マカを含有し、
血中成長ホルモン量上昇作用を有する
ことを特徴としたマカを含有した機能性食品。

【請求項 2】

マカを含有し、
持久力増加作用を有する
ことを特徴としたマカを含有した機能性食品。

【請求項 3】

5 週間以上の継続摂取により、血中成長ホルモン量が上昇する
ことを特徴とした請求項 1 記載のマカを含有した機能性食品。

【請求項 4】

マカのアルコール抽出物を含有している
ことを特徴とする請求項 1 ないし 3 いずれか記載のマカを含有した機能性食品。

【請求項 5】

マカのアルコール抽出物を含有し、
このアルコール抽出物は、60%以上80%以下の含水エタノールを含んでいる
ことを特徴とした請求項 1 ないし 4 いずれか記載のマカを含有した機能性食品。

【発明の詳細な説明】

【0001】

【発明の属する技術分野】

本発明は、マカを含有した機能性食品に関する。

【0002】

【従来の技術】

成長ホルモンは脳下垂体から分泌され、窒素保持能力を改善し、異化能力が低下した場合に脂肪動員と蛋白合成とを促進させる効果が知られている。そして、成長ホルモンは、各種ホルモンによる分泌調節を受け、脈動的な分泌動態があり、日内変動が認められるものの恒常性が保たれている。しかし、加齢などにより成長ホルモンの分泌低下が起こると、シワや白髪の増加、体力および生殖機能の低下をはじめとする、いわゆる老化現象が出やすくなると考えられている。

【0003】

そこで、薬剤ではなく、人体に安全で血中成長ホルモン量上昇作用を有する、いわゆる機能性食品の開発が強く望まれているが、この要望を完全に満たすものは未だないのが現状である。

【0004】

従来、ペルーのアンデス地方の高地原産の根菜であるマカ（レピディウム・メイエニワルプ；*Lepidium meyenii* Walp, *Cruciferae*）が、南米アンデスの原住民の間では、永年伝承薬として不妊治療、滋養強壮などに使われており、民間伝承薬として用いられ、栄養不良、貧血症、不妊症、精力減退症などへの有効性が知られている（例えば、特許文献 1 および 2 参照。）。

【0005】

【特許文献 1】

特開 2000-316528 号公報（第 2-3 頁）

【0006】

【特許文献 2】

特開 2001-136920 号公報（第 5-6 頁）

【0007】

【発明が解決しようとする課題】

しかしながら、上記マカの乾燥物中には、蛋白質や食物繊維含量が比較的高く、特にリジ

10

20

30

40

50

ンやアルギニンの含有量が高く、また特殊成分としてベンジルグルコシレート（ベンジルまたはP-メトキシベンジルイソチオシアネートを生成するグルコシノレート）などのグルコシノレート、ステロイド、フラボノイド、アルカロイド、ポリフェノール、サポニン、テルペン類などの存在が知られているが、これらの成分の生理作用や有効性との関連についてはほとんど知られていないという問題を有している。

【0008】

本発明は、このような点に鑑みなされたもので、血中成長ホルモン量上昇作用を有するマカを含有した機能性食品を提供する。

【0009】

【課題を解決するための手段】

請求項1記載のマカを含有した機能性食品は、マカを含有し、血中成長ホルモン量上昇作用を有するものである。

【0010】

そして、マカを含有した食品を血中成長ホルモン量が低下した場合に摂取することにより、血中成長ホルモン量を適度に上昇できる。

【0011】

請求項2記載のマカを含有した機能性食品は、マカを含有し、持久力増加作用を有するものである。

【0012】

そして、マカを含有した食品を摂取することにより、持久力を増加できる。

【0013】

請求項3記載のマカを含有した機能性食品は、請求項1記載のマカを含有した機能性食品において、5週間以上の継続摂取により、血中成長ホルモン量が上昇するものである。

【0014】

そして、マカを含有した食品を、5週間以上の継続して摂取することにより、いわゆる老化現象の抑制を期待できる。

【0015】

請求項4記載のマカを含有した機能性食品は、請求項1ないし3いずれか記載のマカを含有した機能性食品において、マカのアルコール抽出物を含有しているものである。

【0016】

そして、マカのアルコール抽出物を含有した機能性食品を摂取することにより、マカを含有した食品をより効率良く摂取できる。

【0017】

請求項5記載のマカを含有した機能性食品は、請求項1ないし4いずれか記載のマカを含有した機能性食品において、マカのアルコール抽出物を含有し、このアルコール抽出物は、60%以上80%以下の含水エタノールを含んでいるものである。

【0018】

そして、60%以上80%以下の含水エタノールを含んだマカのアルコール抽出物を含有させることにより、マカを含有した食品をより効率良く摂取できる。

【0019】

【発明の実施の形態】

以下、本発明の実施の一形態のマカを含有した機能性食品を説明する。

【0020】

まず、マカの根の乾燥物（チップ）を30メッシュ以上100メッシュ以下程度に粉碎する。そして、この粉碎したマカの根のチップを、室温（より抽出効率を上げる場合には40℃程度の温度）で48時間以上72時間以下含水アルコールに浸漬してエキスを抽出する。抽出液は、そのまま減圧下に濃縮乾固してもよく、抽出液にデキストリン、ソルビトールなどの乾燥助剤を加えて噴霧乾燥してもよい。また、抽出した乾燥物を再度水に溶解し、活性炭、イオン交換樹脂処理などにより更に精製して濃縮あるいは乾固してもよい。これら乾燥物は、適宜ブレンダーなどで粉碎して使用する。用いられるアルコールは、エ

10

20

30

40

50

タノール、プロパノールなどの低級アルコールであり、通常60%以上80%以下の含水エタノールをアルコールとして含んでいることが好ましい。これらの結果、マカのアルコール抽出エキス乾燥物が得られる。

【0021】

次いで、マカの含水アルコール抽出エキス乾燥物が、これを含有せしめた血中成長ホルモン量上昇作用を有する機能性食品として提供される形態としては、散剤、顆粒、錠剤、糖衣錠、カプセル、液剤、シロップ状のいずれであっても良く、これらは適宜助剤、賦香料とともに賦形されてもよい。用いられる賦形剤、希釈剤としては、ゼラチン、糖類、澱粉類、脂肪酸およびその塩、油脂、タルク、生理食塩水、その他のマスキング剤などが挙げられる。

10

【0022】

これらのものをそのまま服用してもよいが、各種料理品、菓子、キャンデなどの食品に混ぜて服用するのも好都合である。服用量は、個人差、個体差が大きいけれども、通常成人一日当たり、水分8.1%、糖質78.9%を含むアルコール抽出エキス乾燥物として、0.1g以上10g以下である。

【0023】

さらに、マカの乾燥物の含水アルコール抽出乾燥物を含有せしめた血中成長ホルモン量上昇作用を有する機能性食品は、常人が毎日食することに問題はないが、特になんらかの原因で成長ホルモンの分泌が減少し、血中成長ホルモン量が低下したようなときには、少なくとも4週間、通常5週間以上の継続した摂取が望ましい。

20

【0024】

なお、マカの乾燥物中には、蛋白質や食物繊維含量が比較的高く、特にリジンやアルギニンの含有量が高く、また特殊成分としてグルコシノレート、ステロイド、フラボノイド、アルカロイド、ポリフェノール、サポニン、テルペン類などの有機化合物の存在が知られているため、これら特殊成分が活性酸素消去活性、ホルモン様活性、免疫賦活活性などに関係することが推定される。また、これら有機化合物は、特に、ヒトに対して不妊治療に効果のあることからそのグルコシノレートやフラボノイドにフィトエストロゲン作用が予想される。さらに、これら有機化合物により血糖値を低下させる作用が期待できるとともに、美顔や美肌効果なども期待できる。

【0025】

30

【実施例】

以下、本発明のマカのアルコール抽出エキス乾燥物を含有せしめた血中成長ホルモン量上昇作用を有する機能性食品を、実験例を用いて更に詳細に説明する。

【0026】

(実験例1)

マカ・エキスパウダの調製

まず、洗って乾燥させたペルー産マカ（根）（トワ商事株式会社販売）を裁断しチップとし、これを30メッシュ以上40メッシュ以下に粉碎したものを用意した。

【0027】

次いで、98%エタノールに、予め活性炭およびイオン交換樹脂処理した純水を加えて60%に希釈して含水エタノールを調製し、これの600lに、マカの粉碎物120kgを加えて室温で72時間浸漬した。

40

【0028】

そして、固形物を濾去して得た抽出液480l以上490l以下に質量%が3%の食品用デキストリンを添加して混合、これを噴霧乾燥してマカのアルコール抽出噴霧乾燥物（マカ・エキスパウダ）15kgを得、この後、この乾燥物をブレンダーで粉碎して混合した。ここで、このマカ・エキスパウダの分析結果を表1に示す。

【0029】

【表1】

	分析値〔質量％〕
水分	8.1
蛋白質	3.3
脂質	1.2
水不溶性食物繊維	0.0
水可溶性食物繊維	6.7
糖 ※	78.9
灰分	1.8

10

※：計算値

【0030】

(実験例2)

カゼインを蛋白源（20質量％）とするAIN-93組成（リーブスなど、ジャーナル・オブ・ニュートリション、123巻1923頁、1993年（J. Nutrition、123、1923、1993））に準拠した半精製飼料を対照食とし、対照食に実験例1で得たマカ・エキスパウダを質量比で0.5％（0.5M）、5％（5M）添加し、飼料の一般組成が対照食と同一となるように調整したものを実験食（0.5M、5M）とした。これら飼料の組成を表2に示す。なお、AIN-93は、米国国立栄養研究所が1993年に発表したマウスおよびラットを用いた栄養研究のための標準精製飼料組成である。

20

【0031】

【表2】

〔質量％〕

飼料成分	対照食 (MC)	0.5M	5M
マカ・エキスパウダ	0	0.5	5.0
ミルクカゼイン	20.0	20.0	19.8
L・シスチン	0.3	0.3	0.3
大豆油	10.0	9.9	9.9
ミネラル混合物 (AIN-93G)	3.5	3.5	3.5
ビタミン混合物 (AIN-93VX)	1.0	1.0	1.0
セルロースパウダ	5.0	5.0	4.7
コーンスターチ	36.8	36.4	32.8
α-コーンスターチ	13.2	13.2	13.2
シュクロース	10.0	10.0	10.0
コリン酒石酸重水素塩	0.25	0.25	0.25

30

40

【0032】

そして、7週齢の雄性ウイスターラット（日本エスエルシー株式会社より購入）を各群6匹ずつ使用して、1週間対照食にて予備飼育した後、温度22.0±2.0℃、湿度55.0±5.0％、12時間の明暗切り替え方式（明期は午前8時から午後8時まで、暗期は午後8時から午前8時まで）で、飲み水（水道水）とともに、飼料自由摂取下に5週間飼育した。なお、本動物実験は「近畿大学医学部動物実験の指針」に則って実施した。そして、実験の間、1日置きに体重と飼料摂取量とを測定した。結果として、体重測定値を表3に示し、飼料摂取量動向を表4に示す。

【0033】

50

【表 3】

体 重 推 移 (各群 6 匹の平均体重 g)

試験開始 から日目	対照食	0.5 M	5 M	試験開始 から日目	対照食	0.5 M	5 M
2	167	166	166	20	263	260	256
4	183	182	178	22	268	266	262
6	194	197	192	24	273	272	265
8	213	213	208	26	280	276	272
10	223	223	220	28	285	282	275
12	235	236	236	30	288	285	280
14	239	240	238	32	294	292	282
16	244	243	240	34	298	296	289
18	255	255	248	36	304	301	293

【0034】

【表 4】

各群の平均飼料摂取量

〔g/2匹/2日〕

試験開始 から日目	対照食	0.5 M	5 M	試験開始 から日目	対照食	0.5 M	5 M
2	54	50	48	20	64	56	54
4	74	69	62	22	61	60	57
6	70	76	68	24	59	65	57
8	82	79	78	26	58	55	54
10	71	75	77	28	55	57	53
12	66	64	71	30	54	56	55
14	65	64	62	32	54	58	52
16	64	65	56	34	53	54	54
18	63	60	55	36	54	55	55

【0035】

また、実験開始から 2 週間後、無絶食条件でエーテルで麻酔下、鎖骨下静脈からヘパリン（15 単位）含有シリコンコートマイクロスピッツに一部採血（約 1 ml）し、血清を分離（2500 rpm、15 分、4℃）した。また、実験開始より 5 週間後の採血は、無絶食条件でエーテルで麻酔下、腹部大動脈より行い、上記と同一条件の血漿を得た。

【0036】

さらに、実験開始から 2 週間後に得られた血漿、および 5 週間後に得られた血漿中の成長ホルモン量の測定は、ラット成長ホルモン酵素イムノアッセイ（EIA）システム（Amersham Life Science, England）を用いた。そして、2 週間後の血漿中のホルモン量を表 5 に示し、5 週間後に得られた血漿中の成長ホルモン量を表 6 に示す。

【0037】

【表 5】

10

20

30

40

2週間投与後の血漿中成長ホルモン濃度〔ng/ml〕		
対照食	0.5M	5M
92.3±4.9	88.7±8.0	83.6±7.5

n=6、平均値±標準誤差

【0038】

【表6】

10

5週間投与後の血漿中成長ホルモン濃度〔ng/ml〕		
対照食	0.5M	5M
103.2±8.1	120.3±7.8	120.6±12.6

n=6、平均値±標準誤差

【0039】

これらの結果、表3および表4に示すように、マカ・エキスパウダを添加した飼料を投与して飼育したウイスターラットと、対照飼料を投与して飼育したウイスターラットとの成長度（増体量）および飼料摂取量に差は認められず、有害作用も全く認められず、3群とも順調な生育が認められた。

20

【0040】

また、表5および表6に示すように、投与2週間後の血漿中の成長ホルモン量は、実験群（マカを投与した群）において若干低い傾向が観られるが、3群間に有意な差は認められなかった。さらに、投与5週間後に得られた血漿中の成長ホルモン量は、実験群（0.5M、5M両群）で対照群と比較して高い値が認められた。

【0041】

したがって、マカの乾燥物の含水アルコール抽出乾燥物を添加した飼料をウイスターラットに5週間継続して供与して飼育したところ、ラットは順調に生育し、これらウイスターラットにおける血漿中の成長ホルモンが増加した。この結果、主成分として、マカの乾燥物の含水アルコール抽出乾燥物を含有せしめた機能性食品を摂取することにより、人体に安全で、副作用なく、血中成長ホルモン量が上昇することが分かった。

30

【0042】

（実験例3）

次に、本発明のマカのアルコール抽出エキス乾燥物を含有せしめた機能性食品の経口摂取によるマウスの持久力に及ぼす影響を、実験例を用いて説明する。

【0043】

まず、最初に生体機能全体に対する作用を評価する目的で、持久力に対するマカの経口摂取の影響について検討した。

40

【0044】

持久力の評価対象としては、ほぼ成獣に達したと考えられる11週令のDDYマウスを使用した。これらDDYマウスは、研究室に到着した後、1週間馴らし飼育をした後に使用した。また、持久力としては、直径15cmの円筒の内側に各DDYマウスを入れ、これら各DDYマウスを一定速度で走らせて円筒を回転させ、これら各DDYマウスが一定速度で走れなくなるまでの時間（距離）を指標として評価した。なお、走行実験は、一日のうちの同じ時間帯に測定した。

【0045】

また、マカはマカの根の乾燥物を微粉末にした後、水に0.25mg/mlとなるように

50

懸濁し、各 D D Y マウスの胃内にゾンデにて 0. 25 ml を直接単回投与した。

【0046】

ここで、マウスの選抜に関しては、初めどのようなマウスでも走行動態に大きな差はないであろうと仮定して 10 匹のグループに分け測定を始めたが、走行能力あるいは走行意志の欠如などの点で個体差が大きく予備走行テストで選抜しないと単純なグループ分けでは実験に使えないことが判明した。

【0047】

そこで、同一週令のマウスの中で、走行能力に差はあるが常に比較的安定して走行することを基準に、個体をまず選抜した。ところが、平常の走行能力に大きく差のない 2 つのマウス群（対照群および摂取群）を作るのが困難であったため、選抜された常に比較的安定して走行する個体群に対して、マカ投与の前後での走行時間を比較した。

【0048】

次いで、マカの投与容量が食餌の摂取に及ぼす影響に関しては、マカをゾンデで投与する容量 (ml) を変化させてマウスの食餌の摂取量に与える影響を調べた結果、0. 5 ml 以上投与した場合に食餌の摂取量が減少することが判明し、投与容量が 0. 1 ml と決定した。

【0049】

さらに、マカの単回投与がマウスの持久走に与える影響に関しては、上記のように選抜したマウス 7 匹（1 匹死亡）につき、マカの投与前の走行時間（距離）と、0. 1 mg/ml のマカ懸濁液を 0. 25 ml 投与した後の 3 日間走行時間（距離）との変化を測定した。この結果、図 1 ないし図 6 に示すように、6 匹中 4 匹で走行時間が、24 時間後において 25 % 以上 50 % 以下の割合で増加した。なお、後の 2 匹には変化がなかった。また、2 日および 3 日後でも減少加減であるが少し高い割合を示した。さらに、4 日以降は、元の割合に戻ったので、この単回投与量の効果は 2 日以上 3 日以下の間持続すると解釈できた。

【0050】

上述した結果から、マカの経口投与によるマウスの持久力（持久走）に与える影響を検討した。なお、摂取量の正確な量を把握するためゾンデで強制投与する方法を用いて投与した。マウスの食餌摂取量に影響を与えない投与容量 (0. 1 ml) で 1 匹当たり 0. 025 mg/日のマカを与えた結果、6 匹中 4 匹 (66. 6 %) で、持久力の増加が認められ、他の 2 匹においても持久力の低下は示されなかった。

【0051】

また、用いたマウス 10 匹（雄 5 匹、雌 5 匹）中、雌 1 匹が途中死亡し、雄 3 匹が衰弱したのではなく走行状態からの逃避のため測定不能になったため、雌 4 匹および雄 2 匹の結果であるが、図 1 ないし図 6 に示すように、雌では 4 匹中 3 匹 (75 %)、雄では 2 匹中 1 匹 (50 %) の割合で効果が認められた。

【0052】

（実験例 4）

次に、本発明のマカのアルコール抽出エキス乾燥物を含有せしめた機能性食品の連続投与によるマウスの持久力に及ぼす効果を、実験例を用いて説明する。

【0053】

まず、上記実験例 3 と同様の D D Y マウス（雄 2 匹および雌 4 匹）を用意し、これら各マウスそれぞれに上記実験例 3 と同様のマカの乾燥物を複数回数日置きに継続して経口摂取させて連続投与した。

【0054】

具体的には、まず、図 7 ないし図 12 に示すように、実験開始前 3 日間の走行時間を測定し、これら 3 日間における平均走行時間を算出した。なお、これら算出した平均走行時間を、図 7 ないし図 12 中において破線として示した。

【0055】

この後、実験開始日（0 日目）、4 日目、11 日目および 13 日目のそれぞれにおいて各

10

20

30

40

50

マウスにマカの乾燥物をそれぞれ所定量投与した。ここで、図 7 ないし図 12 においては、マカの乾燥物を投与した日を矢印にて示した。

【0056】

この結果、マカの乾燥物を投与させる度に、投与から 1 日後および 2 日後に走行距離が延長したマウスが観察された。具体的には、図 7 ないし図 12 に示すマウス 6 匹中、図 7 に示すマウス（雄 1）、図 9 に示すマウス（雄 3）、図 11 に示すマウス（雌 5）、図 12 に示すマウス（雌 6）の 4 匹のマウスにおいて、マカの乾燥物を投与してから 1 日目および 2 日目に走行距離の延長が確認できた。

【0057】

また、図 10 に示すマウス（雌 4）は、途中（実験開始 6 日目から）で走行距離の測定が不可能となったが、実験開始日でのマカの投与では走行距離の伸びが確認できた。

【0058】

以上のことから、6 匹中 5 匹のマウスにおいて、マカを投与する毎に走行距離の延長が確認できたことにより、マカの摂取により持久力の向上が確認できた。また、図 8 に示すマウス（雌 2）以外の各マウスにおいては、 $P < 0.05$ で統計的に有意差があることが確認できた。

【0059】

（製剤例 1）

実験例 1 で得たマカ・エキスパウダの粉末 1 g に柑橘フレーバ 0.01 g と馬鈴薯澱粉 10 g とを混合し、常法により錠剤（食品）を調製した。

【0060】

（製剤例 2）

実験例 1 で得たマカ・エキスパウダの粉末 0.5 g に、精製白糖 10 g と、柑橘フレーバ 0.05 g とを加えた組成のものに、水を加えて全量 120 ml とし、プラスチックボトル詰めとして飲料製品（ドリンク剤）を調製した。

【0061】

【発明の効果】

請求項 1 記載のマカを含有した機能性食品によれば、マカを含有した機能性食品を摂取することにより、血中成長ホルモン量を上昇できる。

【0062】

請求項 2 記載のマカを含有した機能性食品によれば、マカを含有した機能性食品を摂取することにより、持久力を増加できる。

【0063】

請求項 3 記載のマカを含有した機能性食品によれば、請求項 1 記載のマカを含有した機能性食品の効果に加え、マカを含有した機能性食品を、5 週間以上の継続して摂取することにより、いわゆる老化現象の抑制を期待できる。

【0064】

請求項 4 記載のマカを含有した機能性食品によれば、請求項 1 ないし 3 いずれか記載のマカを含有した機能性食品の効果に加え、マカのアルコール抽出物を含有した機能性食品を摂取することにより、マカを含有した食品をより効率良く摂取できる。

【0065】

請求項 5 記載のマカを含有した機能性食品によれば、請求項 1 ないし 4 いずれか記載のマカを含有した機能性食品の効果に加え、60%以上80%以下の含水エタノールを含んだマカのアルコール抽出物を含有するので、マカを含有した機能性食品をより効率良く摂取できる。

【図面の簡単な説明】

【図 1】本発明のマカを含有した機能性食品をマウス（雄 1）に投与した実験例を示すグラフである。

（a） マカ投与前後の体重の変化を示すグラフ。

（b） マカ投与前後の餌摂取量の変化を示すグラフ。

10

20

30

40

50


- (c) マカ投与前後の運動持久力の変化を示すグラフ。
【図2】同上マカを含有した機能性食品をマウス（雌2）に投与した実験例を示すグラフである。
- (a) マカ投与前後の体重の変化を示すグラフ。
(b) マカ投与前後の餌摂取量の変化を示すグラフ。
(c) マカ投与前後の運動持久力の変化を示すグラフ。
【図3】同上マカを含有した機能性食品をマウス（雄3）に投与した実験例を示すグラフである。
- (a) マカ投与前後の体重の変化を示すグラフ。
(b) マカ投与前後の餌摂取量の変化を示すグラフ。
(c) マカ投与前後の運動持久力の変化を示すグラフ。
【図4】同上マカを含有した機能性食品をマウス（雌4）に投与した実験例を示すグラフである。
- (a) マカ投与前後の体重の変化を示すグラフ。
(b) マカ投与前後の餌摂取量の変化を示すグラフ。
(c) マカ投与前後の運動持久力の変化を示すグラフ。
【図5】同上マカを含有した機能性食品をマウス（雌5）に投与した実験例を示すグラフである。
- (a) マカ投与前後の体重の変化を示すグラフ。
(b) マカ投与前後の餌摂取量の変化を示すグラフ。
(c) マカ投与前後の運動持久力の変化を示すグラフ。
【図6】同上マカを含有した機能性食品をマウス（雌6）に投与した実験例を示すグラフである。
- (a) マカ投与前後の体重の変化を示すグラフ。
(b) マカ投与前後の餌摂取量の変化を示すグラフ。
(c) マカ投与前後の運動持久力の変化を示すグラフ。
【図7】同上マカを含有した機能性食品を連続投与したマウス（雄1）の走行時間を示すグラフである。
- 【図8】同上マカを含有した機能性食品を連続投与したマウス（雌2）の走行時間を示すグラフである。
- 【図9】同上マカを含有した機能性食品を連続投与したマウス（雄3）の走行時間を示すグラフである。
- 【図10】同上マカを含有した機能性食品を連続投与したマウス（雌4）の走行時間を示すグラフである。
- 【図11】同上マカを含有した機能性食品を連続投与したマウス（雌5）の走行時間を示すグラフである。
- 【図12】同上マカを含有した機能性食品を連続投与したマウス（雌6）の走行時間を示すグラフである。

10


20

30


【図 1】




【図 2】




【図 3】




【図 4】




【図 5】



(a)




(b)




(c)


【図 6】



(a)




(b)




(c)


【図 7】




【図 9】




【図 8】




【図 10】



【図 1 1】



【図 1 2】



フロントページの続き

(72)発明者 河村 幸雄

奈良県奈良市朱雀二丁目 3-13

Fターム(参考) 4B018 MD61 ME04

*** NOTICES ***

JPO and NCIP are not responsible for any damages caused by the use of this translation.

1. This document has been translated by computer. So the translation may not reflect the original precisely.
2. **** shows the word which can not be translated.
3. In the drawings, any words are not translated.

CLAIMS

[Claim(s)]

[Claim 1]

MAKA is contained,

It has the amount rise operation of growth hormones in blood.

Functional food containing MAK A characterized by things.

[Claim 2]

MAKA is contained,

It has the increment operation in tenacity.

Functional food containing MAK A characterized by things.

[Claim 3]

By continuation intake for five weeks or more, the amount of growth hormones in blood rises.

Functional food containing MAK A according to claim 1 characterized by things.

[Claim 4]

The alcoholic extract of MAK A is contained.

Claim 1 characterized by things thru/or functional food which contained MAK A of a publication 3 either.

[Claim 5]

The alcoholic extract of MAK A is contained,

This alcoholic extract contains 80% or less of water ethanol 60% or more.

Claim 1 characterized by things thru/or functional food which contained MAK A of a publication 4 either.

[Translation done.]

*** NOTICES ***

JPO and NCIP are not responsible for any damages caused by the use of this translation.

- 1.This document has been translated by computer. So the translation may not reflect the original precisely.
- 2.**** shows the word which can not be translated.
- 3.In the drawings, any words are not translated.

DETAILED DESCRIPTION

[Detailed Description of the Invention]

[0001]

[Field of the Invention]

This invention relates to the functional food containing MAKa.

[0002]

[Description of the Prior Art]

A growth hormone is fossa-hypophysis secreted, nitrogen maintenance capacity is improved, and when catabolism capacity declines, the effectiveness of promoting lipid mobilization and protein synthesis is known. And a growth hormone receives the secretion accommodation by various hormone, and has a pulsation secretion moving state, and homeostasis is maintained although a daily variation is accepted. However, if the secretion fall of a growth hormone takes place by aging etc., it is thought that it becomes easy to come out of the so-called degraded phenomena including the fall of the increment in Siwa or canities, physical strength, and the generative function.

[0003]

Then, although it is safe for the body instead of drugs and development of the so-called functional food which has the amount rise operation of growth hormones in blood is desired strongly, the present condition is that there is still nothing that fills this request completely.

[0004]

Conventionally, MAKa (repi DIUMU MEIENIWARUPU; *Lepidium meyenii*-Walp, Cruciferae) which is the high-ground native root vegetables of the Andes district in Peru is used for infertility treatment, nourishment sthenia, etc. as a tradition medicine for years among the natives of the South America Andes, it is used as a folklore medicine, and the effectiveness to malnutrition, anemia, infertility, energy *****, etc. is known (for example, the patent reference 1 and 2 reference.).

[0005]

[Patent reference 1]

JP,2000-316528,A (the two - 3rd page)

[0006]

[Patent reference 2]

JP,2001-136920,A (the five - 6th page)

[0007]

[Problem(s) to be Solved by the Invention]

However, in the dry matter of above-mentioned MAKa, protein and a dietary fiber content are comparatively high. especially -- the content of a lysine or an arginine -- high -- moreover -- as a special component -- a benzyl guru -- chewiness -- glucosinolates, such as a rate (glucosinolate which generates benzyl or P-methoxyl benzyl isothiocyanate), -- Although existence of a steroid, flavonoid, alkaloid, polyphenol, a saponin, terpenes, etc. is known, about the physiological function of these components, or relation with effectiveness, it has the problem that it is hardly known.

[0008]

This invention was made in view of such a point, and offers the functional food containing MAKa which has the amount rise operation of growth hormones in blood.

[0009]

[Means for Solving the Problem]

The functional food containing MAKa according to claim 1 contains MAKa, and has the amount rise operation of growth hormones in blood.

[0010]

And the amount of growth hormones in blood can be moderately gone up by taking in the food containing MAKa, when the amount of growth hormones in blood falls.

[0011]

The functional food containing MAKa according to claim 2 contains MAKa, and has the increment operation in tenacity.

[0012]

And tenacity can be increased by taking in the food containing MAKa.

[0013]

In the functional food with which the functional food containing MAKa according to claim 3 contained MAKa according to claim 1, the amount of growth hormones in blood rises by continuation intake for five weeks or more.

[0014]

And the so-called control of a degraded phenomenon is [the food containing MAKa] expectable by [for five weeks or more] taking in continuously.

[0015]

The functional food containing MAKa according to claim 4 contains the alcoholic extract of MAKa in claim 1 thru/or the functional food which contained MAKa of a publication 3 either.

[0016]

And the food containing MAKa can be more efficiently taken in by taking in the functional food containing the alcoholic extract of MAKa.

[0017]

The functional food containing MAKa according to claim 5 contains the alcoholic extract of MAKa in claim 1 thru/or the functional food which contained MAKa of a publication 4 either, and this alcoholic extract contains 80% or less of water ethanol 60% or more.

[0018]

And the food containing MAKa can be more efficiently taken in by making the alcoholic extract of MAKa which contained 80% or less of water ethanol 60% or more contain.

[0019]

[Embodiment of the Invention]

Hereafter, the functional food containing MAKa of one gestalt of operation of this invention is explained.

[0020]

First, the dry matter (chip) of the root of MAKa is ground to 30 or more mesh 100 or less mesh extent. And the chip of the root of this ground MAKa is immersed in water alcohol for 72 or less hours for 48 hours or more at a room temperature (when gathering extraction efficiency, it is the temperature of about 40 degrees C), and an extractive matter is extracted. An extract may carry out concentration hardening by drying under reduced pressure as it is, and may add and carry out spray drying of the drying aid, such as a dextrin and a sorbitol, to an extract. Moreover, the extracted dry matter is again dissolved in water, and activated carbon, ion-exchange-resin processing, etc. may refine further, and you may condense or harden by drying. These dry matters are suitably ground and used with a blender etc. The alcohol used is lower alcohol, such as ethanol and propanol, and it is desirable that 80% or less of water ethanol is usually included as alcohol 60% or more. The alcoholic extract extractives dry matter of MAKa is obtained these results.

[0021]

Subsequently, the water alcoholic extract extractives dry matters of MAKa may be powder, granulation, a tablet, a sugar-coated tablet, a capsule, liquids and solutions, and sirupy any as a gestalt offered as functional food which has the amount rise operation of growth hormones in blood which made this contain, and size enlargement of these may be suitably carried out with an assistant and the charge of aromatizing. As the excipient used and a diluent, the masking reagent of gelatin, a saccharide, starch, a fatty acid and its salt, fats and oils, talc, a physiological saline, and others etc. is mentioned.

[0022]

Although these things may be taken as they are, it is also convenient to mix food, such as various cooking articles, confectionery, and Kandy, and to take. Although individual difference and individual difference of a dose

are large, it is 0.1g or more 10g or less as an alcoholic extract extractives dry matter which usually contains per adult day, 8.1% of moisture, and 78.9% of sugar.

[0023]

Furthermore, although it is satisfactory for usual state people eating the functional food which has the amount rise operation of growth hormones in blood which made the water alcoholic extract dry matter of the dry matter of MAKKA contain every day, when secretion of a growth hormone decreases especially by a certain cause and the amount of growth hormones in blood falls, the continued intake for five weeks or more is usually desirable for at least four weeks.

[0024]

In addition, in the dry matter of MAKKA, protein and a dietary fiber content are comparatively high, especially the content of a lysine or an arginine is high, and since existence of organic compounds, such as glucosinolate, a steroid, flavonoid, alkaloid, polyphenol, a saponin, and terpenes, is known as a special component, it is presumed that these special components are related to active oxygen elimination activity, hormone Mr. activity, immunity activation activity, etc. Moreover, since these organic compounds have effectiveness in infertility treatment to Homo sapiens especially, a phytoestrogen operation is expected by the glucosinolate and flavonoid. Furthermore, while the operation which reduces the blood sugar level with these organic compounds is expectable, facial treatment, a beautiful skin effect, etc. are expectable.

[0025]

[Example]

The functional food which has hereafter the amount rise operation of growth hormones in blood which made the alcoholic extract extractives dry matter of MAKKA of this invention contain is further explained to a detail using the example of an experiment.

[0026]

(Example 1 of an experiment)

Preparation of MAKKA extractives powder

First, it washed, dried MAKKA from Peru (root) (TOWA business-affairs incorporated company sale) was judged, it considered as the chip, and what ground this to 30 or more-mesh 40 or less meshes was prepared.

[0027]

Subsequently, activated carbon and the pure water which carried out ion-exchange-resin processing were beforehand added to ethanol 98%, it diluted to 60%, and water ethanol was prepared, and 120kg of grinding objects of MAKKA was added to 600l. of this, and it was immersed in it at the room temperature for 72 hours.

[0028]

And mass % added 3% of food-grade dextrin to 490l. or less of 480l. or more extracts which filtered out and obtained the solid, and was mixed to them, and spray drying of this was carried out and 15kg (MAKKA extractives powder) of alcoholic extract spray drying objects of MAKKA was obtained, and after this, the blender ground this dry matter and it mixed. Here, the analysis result of this MAKKA extractives powder is shown in Table 1.

[0029]

[Table 1]

	分析値〔質量%〕
水分	8.1
蛋白質	3.3
脂質	1.2
水不溶性食物繊維	0.0
水可溶性食物繊維	6.7
糖 ※	78.9
灰分	1.8

※：計算値

[0030]

(Example 2 of an experiment)

The AIN-93 presentation which makes casein the source of protein (20 mass %) (Reeves etc.) Journal OBU new TORISHON, 123-volume 1923 pages, 1993 (J. it Nutrition(s)) Half-purified diet based on 123, 1923, and 1993 is considered as contrast foods. Addition of the MAKKA extractives powder obtained in the example 1 of an experiment to contrast foods was carried out 5% (5M) 0.5% (0.5M) with the mass ratio, and what was adjusted so that the proximate composition of feed might become the same as that of contrast foods was made into experimental diet (0.5M, 5M). The presentation of these feed is shown in Table 2. In addition, AIN-93 are the standard purified diet presentation for the nutrition research using the mouse and rat which the U.S. National Institute of Nutrition announced in 1993.

[0031]

[Table 2]

〔質量％〕

飼 料 成 分	対照食 (MC)	0.5 M	5 M
マカ・エキスパウダ	0	0.5	5.0
ミルクカゼイン	20.0	20.0	19.8
L・シスチン	0.3	0.3	0.3
大豆油	10.0	9.9	9.9
ミネラル混合物 (A I N - 9 3 G)	3.5	3.5	3.5
ビタミン混合物 (A I N - 9 3 V X)	1.0	1.0	1.0
セルロースパウダ	5.0	5.0	4.7
コーンスターチ	36.8	36.4	32.8
α-コーンスターチ	13.2	13.2	13.2
シュクロース	10.0	10.0	10.0
コリン酒石酸重水素塩	0.25	0.25	0.25

[0032]

And after using each six groups at a time 7-weeks old male Wistar rats (it purchases from Japan SLC, Inc.) and carrying out preliminary breeding by contrast foods for one week, it is the temperature of 22.0**2.0 degrees C, the humidity of 55.0**5.0%, and the light-and-darkness change method (for a ** term, a dark term is from 8:00 p.m. to 8:00 a.m. from 8:00 a.m. to 8:00 p.m.) of 12 hours, and bred for five weeks under feed free intake with drinking water (tap water). In addition, this animal experiment was carried out in conformity with "the guide of the Kinki University medical department animal experiment." And weight and the amount of feed intake were measured every other day during the experiment. As a result, weight measured value is shown in Table 3, and the amount trend of feed intake is shown in Table 4.

[0033]

[Table 3]

体 重 推 移 (各群6匹の平均体重 g)

試験開始 から日目	対照食	0.5 M	5 M	試験開始 から日目	対照食	0.5 M	5 M
2	167	166	166	20	263	260	256
4	183	182	178	22	268	266	262
6	194	197	192	24	273	272	265
8	213	213	208	26	280	276	272
10	223	223	220	28	285	282	275
12	235	236	236	30	288	285	280
14	239	240	238	32	294	292	282
16	244	243	240	34	298	296	289
18	255	255	248	36	304	301	293

[0034]

[Table 4]

各群の平均飼料摂取量

[g/2匹/2日]

試験開始 から日目	対照食	0.5 M	5 M	試験開始 から日目	対照食	0.5 M	5 M
2	54	50	48	20	64	56	54
4	74	69	62	22	61	60	57
6	70	76	68	24	59	65	57
8	82	79	78	26	58	55	54
10	71	75	77	28	55	57	53
12	66	64	71	30	54	56	55
14	65	64	62	32	54	58	52
16	64	65	56	34	53	54	54
18	63	60	55	36	54	55	55

[0035]

Moreover, it collected blood from the subclavian vein in part to the heparin (15 units) content silicon coat micro spitz under anesthesia with the ether by the conditions of not abstaining from food, after [of experiment initiation] two weeks (about 1ml), and the blood serum was separated (2500rpm, 15 minutes, 4 degrees C). Moreover, blood collecting five weeks after was performed on the conditions of not abstaining from food, from experiment initiation, the ether performed it from the abdominal aorta under anesthesia, and the plasma of the same conditions as the above was obtained.

[0036]

Furthermore, measurement of the amount of growth hormones in the plasma obtained after [of experiment initiation] two weeks and the plasma obtained after five weeks used the rat growth hormone enzyme immunoassay (EIA) system (Amersham Life Science, England). And the amount of hormone in the plasma of two weeks after is shown in Table 5, and the amount of growth hormones in the plasma obtained after five weeks is shown in Table 6.

[0037]

[Table 5]

2 週間投与後の血漿中成長ホルモン濃度〔ng/ml〕		
対照食	0.5 M	5 M
92.3 ± 4.9	88.7 ± 8.0	83.6 ± 7.5

n=6、平均値±標準誤差

[0038]
[Table 6]

5 週間投与後の血漿中成長ホルモン濃度〔ng/ml〕		
対照食	0.5 M	5 M
103.2 ± 8.1	120.3 ± 7.8	120.6 ± 12.6

n=6、平均値±標準誤差

[0039]
As shown in Table 3 and 4 these results, the difference was not accepted, but adverse reaction was not accepted in whenever [growth-with Wistar-rats / which prescribed for the patient and bred the feed which added MAKa extractives powder /, and Wistar rats which prescribed for the patient and bred contrast feed], and (gain of body weight), the amount of feed intake at all, either, but growth with all three favorable groups was accepted in them.

[0040]
Moreover, although, as for the amount of growth hormones in the plasma after two weeks of administration, the inclination low a little was seen in the experimental group (group which prescribed MAKa for the patient) as shown in Table 5 and 6, the significant difference was not accepted in 3 between groups. Furthermore, as for the amount of growth hormones in the plasma obtained after five weeks of administration, as compared with the control group, the high value was accepted by the experimental group (0.5M, 5 M car group).

[0041]
Therefore, when it continued for five weeks to Wistar rats, they were supplied with the feed which added the water alcoholic extract dry matter of the dry matter of MAKa and it was bred to them, the rat was grown favorably and the growth hormone in the plasma in these Wistar rats increased. Consequently, it is safe for the body, there is no side effect, and by taking in the functional food which made the water alcoholic extract dry matter of the dry matter of MAKa contain as a principal component showed that the amount of growth hormones in blood rose.

[0042]
(Example 3 of an experiment)
Next, the effect affect the tenacity of the mouse by the ingestion of functional food which made the alcoholic extract extractives dry matter of MAKa of this invention contain is explained using the example of an experiment.

[0043]
First, the purpose which evaluates the operation over the whole living body function to the beginning considered the effect of the ingestion of MAKa to tenacity.

[0044]
The DDY mouse of 11 weeks old considered to have reached the adult mostly as a candidate for evaluation of tenacity was used. These DDY(s) mouse was used after arriving at a laboratory, and carrying out training breeding for one week. Moreover, as tenacity, each DDY mouse was put in inside the cylinder with a diameter of

15cm, each [these] DDY mouse was run with constant speed, the cylinder was rotated, and time amount (distance) until it becomes impossible for each [these] DDY mouse to run with constant speed was evaluated as an index. In addition, the transit experiment was measured in the same time zone of the days.

[0045]

Moreover, after MAKKA used the dry matter of the root of MAKKA as impalpable powder, it was suspended so that it might become water in ml and 0.25mg /, and carried out direct single-dose administration of the 0.25ml by the sound into the stomach of each DDY mouse.

[0046]

Here, although it divided into the group of ten animals, having assumed whether a big difference would have any mice in a transit moving state and measurement was begun at first about selection of a mouse, when individual difference did not select by the preliminary transit test greatly in respect of lack of transit capacity or transit volition etc., it became clear by the simple group division that it could not use for an experiment.

[0047]

Then, the individual was first selected on the basis of being stabilized comparatively and always running in the mouse of the same weeks old, although a difference is in transit capacity. However, since it was difficult to make two mouse groups (a control group and intake group) which do not have a difference in normal transit capacity greatly, it was stabilized comparatively and the transit time in MAKKA administration order was always compared to the selected population it runs.

[0048]

Subsequently, as a result of investigating the effect which the capacity (ml) which prescribes MAKKA for the patient by the sound is changed about the effect the administration capacity of MAKKA affects intake of a diet, and it has on the intake of the diet of a mouse, when 0.5ml or more was prescribed for the patient, it became clear that the intake of a diet decreased and it was decided that administration capacity would be 0.1ml.

[0049]

Furthermore, change with the transit time before administration of MAKKA (distance) and the transit time during three days (distance) after prescribing 0.25ml of 0.1mg [/ml] MAKKA suspension for the patient was measured about seven mice (one-animal death) which the single-dose administration of MAKKA selected as mentioned above about the effect which it has on the endurance running of a mouse. consequently, it is shown in drawing 1 thru/or drawing 6 -- as -- the inside of six animals -- four animals -- the transit time -- 24 hours after -- setting -- coming out -- 50% or less 25% or more -- it came out comparatively and increased. In addition, there was no change in two next animals. Moreover, although it was a reduction degree also after two days and three days, the somewhat high rate was shown. Furthermore, since it returned to the original rate after the 4th, it has been interpreted as maintaining the effectiveness of this amount of single-dose administration during the 3rd or less the 2nd day or more.

[0050]

From the result mentioned above, the effect which it has on the tenacity (endurance running) of the mouse by internal use of MAKKA was considered. In addition, in order to grasp an amount with exact intake, a medicine was prescribed for the patient using the approach of carrying out forcible administration by the sound. As a result of giving 0.025mg [per animal / /] MAKKA day by the administration capacity (0.1ml) which does not affect the diet intake of a mouse, among six animals, by four animals (66.6%), the increment in tenacity was accepted and the fall of tenacity was not shown in other two animals.

[0051]

Moreover, although it is as a result of four females and two males since one female died the middle among ten used mice (five males, five females), and three males did not become weak but it became measurement impossible for escape from a run state As shown in drawing 1 thru/or drawing 6 , among four animals, for the female, effectiveness was accepted by three animals (75%), and was accepted at a rate of one animal (50%) among two animals by the male.

[0052]

(Example 4 of an experiment)

Next, the effectiveness exerted on the tenacity of the mouse by the repetitive administration of the functional food which made the alcoholic extract extractives dry matter of MAKKA of this invention contain is explained using the example of an experiment.

[0053]

First, the same DDY mouse (two males and four females) as the above-mentioned example 3 of an experiment

was prepared, the ingestion of the dry matter of the same MAKa as the above-mentioned example 3 of an experiment was continued and carried out to each of each [these] mouse every count day of plurality, and repetitive administration was carried out.

[0054]

First, as shown in drawing 7 thru/or drawing 12 , the transit time for front [experiment initiation] three days was measured, and, specifically, the mean transit time for these three days was computed. In addition, the these-computed mean transit time was shown as a broken line in drawing 7 thru/or drawing 12 .

[0055]

Then, in each on the 11th and the 13th, specified quantity administration of the dry matter of MAKa was carried out [the experiment opening day (the 0th day) and] on the 4th at each mouse, respectively. Here, in drawing 7 thru/or drawing 12 , the arrow head showed the day which prescribed the dry matter of MAKa for the patient.

[0056]

Consequently, the mouse which mileage extended one day and two days after administration whenever it made the dry matter of MAKa prescribe for the patient was observed. Extension of mileage has been checked on the 1st and the 2nd after prescribing the dry matter of MAKa for the patient in four mice, the mouse (male 1) specifically shown in the inside of six mice shown in drawing 7 thru/or drawing 12 , and drawing 7 , the mouse (male 3) shown in drawing 9 , the mouse (female 5) shown in drawing 11 , and the mouse (female 6) shown in drawing 12 .

[0057]

Moreover, although the measurement of mileage of the mouse (female 4) shown in drawing 10 became impossible from the 6th day of experiment initiation on the way, it has checked stretch of mileage in administration of MAKa in an experiment opening day.

[0058]

From the above thing, among six animals, whenever it prescribed MAKa for the patient in five mice, improvement in tenacity has been checked by intake of MAKa by having checked extension of mileage. Moreover, in each mouse other than the mouse (female 2) shown in drawing 8 , it has checked that there was a significant difference statistically by $P < 0.05$.

[0059]

(Example 1 of pharmaceutical preparation)

Citrus flavor 0.01g and 10g of potato starch were mixed to 1g of powder of the MAKa extractives powder obtained in the example 1 of an experiment, and the tablet (food) was prepared with the conventional method.

[0060]

(Example 2 of pharmaceutical preparation)

Water was added to the thing of a presentation which added 10g of purified sucrose, and citrus flavor 0.05g to 0.5g of **** of the MAKa extractives powder obtained in the example 1 of an experiment, it considered as 120ml of whole quantity, and the drink product (drinkable preparations) was prepared as plastics bottle stuffing.

[0061]

[Effect of the Invention]

According to the functional food containing MAKa according to claim 1, the amount of growth hormones in blood can be gone up by taking in the functional food containing MAKa.

[0062]

According to the functional food containing MAKa according to claim 2, tenacity can be increased by taking in the functional food containing MAKa.

[0063]

According to the functional food containing MAKa according to claim 3, in addition to the effectiveness of the functional food containing MAKa according to claim 1, the so-called control of a degraded phenomenon is [the functional food containing MAKa] expectable by [for five weeks or more] taking in continuously.

[0064]

According to the functional food containing MAKa according to claim 4, the food containing MAKa can be more efficiently taken in by taking in the functional food containing the alcoholic extract of MAKa in addition to claim 1 thru/or the effectiveness of functional food which contained MAKa of a publication 3 either.

[0065]

Since the alcoholic extract of MAKa which contained 80% or less of water ethanol 60% or more is contained [according to the functional food containing MAKa according to claim 5] in addition to claim 1 thru/or the

effectiveness of functional food which contained MAKa of a publication 4 either, the functional food containing MAKa can be taken in more efficiently.

[Brief Description of the Drawings]

[Drawing 1] It is the graph which shows the example of an experiment which medicated the mouse (male 1) with the functional food containing MAKa of this invention.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 2] It is the graph which shows the example of an experiment which medicated the mouse (female 2) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 3] It is the graph which shows the example of an experiment which medicated the mouse (male 3) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 4] It is the graph which shows the example of an experiment which medicated the mouse (female 4) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 5] It is the graph which shows the example of an experiment which medicated the mouse (female 5) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 6] It is the graph which shows the example of an experiment which medicated the mouse (female 6) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 7] It is the graph which shows the transit time of the mouse (male 1) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 8] It is the graph which shows the transit time of the mouse (female 2) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 9] It is the graph which shows the transit time of the mouse (male 3) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 10] It is the graph which shows the transit time of the mouse (female 4) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 11] It is the graph which shows the transit time of the mouse (female 5) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 12] It is the graph which shows the transit time of the mouse (female 6) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Translation done.]

*** NOTICES ***

JP0 and NCIP1 are not responsible for any damages caused by the use of this translation.

- 1.This document has been translated by computer. So the translation may not reflect the original precisely.
- 2.**** shows the word which can not be translated.
- 3.In the drawings, any words are not translated.

TECHNICAL FIELD

[Field of the Invention]

This invention relates to the functional food containing MAKKA.

[0002]

[Translation done.]

*** NOTICES ***

JPO and NCIP are not responsible for any damages caused by the use of this translation.

- 1.This document has been translated by computer. So the translation may not reflect the original precisely.
- 2.**** shows the word which can not be translated.
- 3.In the drawings, any words are not translated.

PRIOR ART

[Description of the Prior Art]

A growth hormone is fossa-hypophysis secreted, nitrogen maintenance capacity is improved, and when catabolism capacity declines, the effectiveness of promoting lipid mobilization and protein synthesis is known. And a growth hormone receives the secretion accommodation by various hormone, and has a pulsation secretion moving state, and homeostasis is maintained although a daily variation is accepted. However, if the secretion fall of a growth hormone takes place by aging etc., it is thought that it becomes easy to come out of the so-called degraded phenomena including the fall of the increment in Siwa or canities, physical strength, and the generative function.

[0003]

Then, although it is safe for the body instead of drugs and development of the so-called functional food which has the amount rise operation of growth hormones in blood is desired strongly, the present condition is that there is still nothing that fills this request completely.

[0004]

Conventionally, MAKU (repi DIUMU MEIENIWARUPU; *Lepidium meyenii*-Walp, Cruciferae) which is the high-ground native root vegetables of the Andes district in Peru is used for infertility treatment, nourishment sthenia, etc. as a tradition medicine for years among the natives of the South America Andes, it is used as a folklore medicine, and the effectiveness to malnutrition, anemia, infertility, energy *****, etc. is known (for example, the patent reference 1 and 2 reference.).

[0005]

[Patent reference 1]

JP,2000-316528,A (the two - 3rd page)

[0006]

[Patent reference 2]

JP,2001-136920,A (the five - 6th page)

[0007]

[Translation done.]

*** NOTICES ***

JPO and NCIP are not responsible for any damages caused by the use of this translation.

1. This document has been translated by computer. So the translation may not reflect the original precisely.
2. **** shows the word which can not be translated.
3. In the drawings, any words are not translated.

EFFECT OF THE INVENTION

[Effect of the Invention]

According to the functional food containing MAKa according to claim 1, the amount of growth hormones in blood can be gone up by taking in the functional food containing MAKa.

[0062]

According to the functional food containing MAKa according to claim 2, tenacity can be increased by taking in the functional food containing MAKa.

[0063]

According to the functional food containing MAKa according to claim 3, in addition to the effectiveness of the functional food containing MAKa according to claim 1, the so-called control of a degraded phenomenon is [the functional food containing MAKa] expectable by [for five weeks or more] taking in continuously.

[0064]

According to the functional food containing MAKa according to claim 4, the food containing MAKa can be more efficiently taken in by taking in the functional food containing the alcoholic extract of MAKa in addition to claim 1 thru/or the effectiveness of functional food which contained MAKa of a publication 3 either.

[0065]

Since the alcoholic extract of MAKa which contained 80% or less of water ethanol 60% or more is contained [according to the functional food containing MAKa according to claim 5] in addition to claim 1 thru/or the effectiveness of functional food which contained MAKa of a publication 4 either, the functional food containing MAKa can be taken in more efficiently.

[Translation done.]

*** NOTICES ***

JPO and NCIP are not responsible for any damages caused by the use of this translation.

- 1.This document has been translated by computer. So the translation may not reflect the original precisely.
- 2.**** shows the word which can not be translated.
- 3.In the drawings, any words are not translated.

TECHNICAL PROBLEM

[Problem(s) to be Solved by the Invention]

However, in the dry matter of above-mentioned MAKAs, protein and a dietary fiber content are comparatively high. especially -- the content of a lysine or an arginine -- high -- moreover -- as a special component -- a benzyl group -- chewiness -- glucosinolates, such as a rate (glucosinolate which generates benzyl or P-methoxyl benzyl isothiocyanate), -- Although existence of a steroid, flavonoid, alkaloid, polyphenol, a saponin, terpenes, etc. is known, about the physiological function of these components, or relation with effectiveness, it has the problem that it is hardly known.

[0008]

This invention was made in view of such a point, and offers the functional food containing MAKAs which has the amount rise operation of growth hormones in blood.

[0009]

[Translation done.]

*** NOTICES ***

JPO and NCIP are not responsible for any damages caused by the use of this translation.

1. This document has been translated by computer. So the translation may not reflect the original precisely.
2. **** shows the word which can not be translated.
3. In the drawings, any words are not translated.

MEANS

[Means for Solving the Problem]

The functional food containing MAKa according to claim 1 contains MAKa, and has the amount rise operation of growth hormones in blood.

[0010]

And the amount of growth hormones in blood can be moderately gone up by taking in the food containing MAKa, when the amount of growth hormones in blood falls.

[0011]

The functional food containing MAKa according to claim 2 contains MAKa, and has the increment operation in tenacity.

[0012]

And tenacity can be increased by taking in the food containing MAKa.

[0013]

In the functional food with which the functional food containing MAKa according to claim 3 contained MAKa according to claim 1, the amount of growth hormones in blood rises by continuation intake for five weeks or more.

[0014]

And the so-called control of a degraded phenomenon is [the food containing MAKa] expectable by [for five weeks or more] taking in continuously.

[0015]

The functional food containing MAKa according to claim 4 contains the alcoholic extract of MAKa in claim 1 thru/or the functional food which contained MAKa of a publication 3 either.

[0016]

And the food containing MAKa can be more efficiently taken in by taking in the functional food containing the alcoholic extract of MAKa.

[0017]

The functional food containing MAKa according to claim 5 contains the alcoholic extract of MAKa in claim 1 thru/or the functional food which contained MAKa of a publication 4 either, and this alcoholic extract contains 80% or less of water ethanol 60% or more.

[0018]

And the food containing MAKa can be more efficiently taken in by making the alcoholic extract of MAKa which contained 80% or less of water ethanol 60% or more contain.

[0019]

[Embodiment of the Invention]

Hereafter, the functional food containing MAKa of one gestalt of operation of this invention is explained.

[0020]

First, the dry matter (chip) of the root of MAKa is ground to 30 or more mesh 100 or less mesh extent. And the chip of the root of this ground MAKa is immersed in water alcohol for 72 or less hours for 48 hours or more at a room temperature (when gathering extraction efficiency, it is the temperature of about 40 degrees C), and an extractive matter is extracted. An extract may carry out concentration hardening by drying under reduced pressure as it is, and may add and carry out spray drying of the drying aid, such as a dextrin and a sorbitol, to an extract. Moreover, the extracted dry matter is again dissolved in water, and activated carbon, ion-exchange-resin processing, etc. may refine further, and you may condense or harden by drying. These dry matters are

suitably ground and used with a blender etc. The alcohol used is lower alcohol, such as ethanol and propanol, and it is desirable that 80% or less of water ethanol is usually included as alcohol 60% or more. The alcoholic extract extractives dry matter of MAKKA is obtained these results.

[0021]

Subsequently, the water alcoholic extract extractives dry matters of MAKKA may be powder, granulation, a tablet, a sugar-coated tablet, a capsule, liquids and solutions, and sirupy any as a gestalt offered as functional food which has the amount rise operation of growth hormones in blood which made this contain, and size enlargement of these may be suitably carried out with an assistant and the charge of aromatizing. As the excipient used and a diluent, the masking reagent of gelatin, a saccharide, starch, a fatty acid and its salt, fats and oils, talc, a physiological saline, and others etc. is mentioned.

[0022]

Although these things may be taken as they are, it is also convenient to mix food, such as various cooking articles, confectionery, and Kandy, and to take. Although individual difference and individual difference of a dose are large, it is 0.1g or more 10g or less as an alcoholic extract extractives dry matter which usually contains per adult day, 8.1% of moisture, and 78.9% of sugar.

[0023]

Furthermore, although it is satisfactory for usual state people eating the functional food which has the amount rise operation of growth hormones in blood which made the water alcoholic extract dry matter of the dry matter of MAKKA contain every day, when secretion of a growth hormone decreases especially by a certain cause and the amount of growth hormones in blood falls, the continued intake for five weeks or more is usually desirable for at least four weeks.

[0024]

In addition, in the dry matter of MAKKA, protein and a dietary fiber content are comparatively high, especially the content of a lysine or an arginine is high, and since existence of organic compounds, such as glucosinolate, a steroid, flavonoid, alkaloid, polyphenol, a saponin, and terpenes, is known as a special component, it is presumed that these special components are related to active oxygen elimination activity, hormone Mr. activity, immunity activation activity, etc. Moreover, since these organic compounds have effectiveness in infertility treatment to Homo sapiens especially, a phytoestrogen operation is expected by the glucosinolate and flavonoid. Furthermore, while the operation which reduces the blood sugar level with these organic compounds is expectable, facial treatment, a beautiful skin effect, etc. are expectable.

[0025]

[Translation done.]

* NOTICES *

JPO and NCIP are not responsible for any damages caused by the use of this translation.

- 1.This document has been translated by computer. So the translation may not reflect the original precisely.
- 2.**** shows the word which can not be translated.
- 3.In the drawings, any words are not translated.

EXAMPLE

[Example]

The functional food which has hereafter the amount rise operation of growth hormones in blood which made the alcoholic extract extractives dry matter of MAKAKA of this invention contain is further explained to a detail using the example of an experiment.

[0026]

(Example 1 of an experiment)

Preparation of MAKAKA extractives powder

First, it washed, dried MAKAKA from Peru (root) (TOWA business-affairs incorporated company sale) was judged, it considered as the chip, and what ground this to 30 or more-mesh 40 or less meshes was prepared.

[0027]

Subsequently, activated carbon and the pure water which carried out ion-exchange-resin processing were beforehand added to ethanol 98%, it diluted to 60%, and water ethanol was prepared, and 120kg of grinding objects of MAKAKA was added to 600l. of this, and it was immersed in it at the room temperature for 72 hours.

[0028]

And mass % added 3% of food-grade dextrin to 490l. or less of 480l. or more extracts which filtered out and obtained the solid, and was mixed to them, and spray drying of this was carried out and 15kg (MAKAKA extractives powder) of alcoholic extract spray drying objects of MAKAKA was obtained, and after this, the blender ground this dry matter and it mixed. Here, the analysis result of this MAKAKA extractives powder is shown in Table 1.

[0029]

[Table 1]

	分析値〔質量%〕
水分	8 . 1
蛋白質	3 . 3
脂質	1 . 2
水不溶性食物纖維	0 . 0
水可溶性食物纖維	6 . 7
糖 ※	7 8 . 9
灰分	1 . 8

※ : 計算値

[0030]

(Example 2 of an experiment)

The AIN-93 presentation which makes casein the source of protein (20 mass %) (Reeves etc.) Journal OBU new TORISHON, 123-volume 1923 pages, 1993 (J. it Nutrition(s)) Half-purified diet based on 123, 1923, and 1993 is considered as contrast foods. Addition of the MAKAKA extractives powder obtained in the example 1 of an experiment to contrast foods was carried out 5% (5M) 0.5% (0.5M) with the mass ratio, and what was adjusted so

that the proximate composition of feed might become the same as that of contrast foods was made into experimental diet (0.5M, 5M). The presentation of these feed is shown in Table 2. In addition, AIN-93 are the standard purified diet presentation for the nutrition research using the mouse and rat which the U.S. National Institute of Nutrition announced in 1993.

[0031]

[Table 2]

〔質量％〕

飼 料 成 分	対照食 (MC)	0.5 M	5 M
マカ・エキスパウダ	0	0.5	5.0
ミルクカゼイン	20.0	20.0	19.8
L・シスチン	0.3	0.3	0.3
大豆油	10.0	9.9	9.9
ミネラル混合物 (A I N-93G)	3.5	3.5	3.5
ビタミン混合物 (A I N-93VX)	1.0	1.0	1.0
セルロースパウダ	5.0	5.0	4.7
コーンスターチ	36.8	36.4	32.8
α-コーンスターチ	13.2	13.2	13.2
シュクロース	10.0	10.0	10.0
コリン酒石酸重水素塩	0.25	0.25	0.25

[0032]

And after using each six groups at a time 7-weeks old male Wistar rats (it purchases from Japan SLC, Inc.) and carrying out preliminary breeding by contrast foods for one week, it is the temperature of 22.0**2.0 degrees C, the humidity of 55.0**5.0%, and the light-and-darkness change method (for a ** term, a dark term is from 8:00 p.m. to 8:00 a.m. from 8:00 a.m. to 8:00 p.m.) of 12 hours, and bred for five weeks under feed free intake with drinking water (tap water). In addition, this animal experiment was carried out in conformity with "the guide of the Kinki University medical department animal experiment." And weight and the amount of feed intake were measured every other day during the experiment. As a result, weight measured value is shown in Table 3, and the amount trend of feed intake is shown in Table 4.

[0033]

[Table 3]

体 重 推 移 (各群 6 匹の平均体重 g)

試験開始 から日目	対照食	0.5 M	5 M	試験開始 から日目	対照食	0.5 M	5 M
2	1 6 7	1 6 6	1 6 6	2 0	2 6 3	2 6 0	2 5 6
4	1 8 3	1 8 2	1 7 8	2 2	2 6 8	2 6 6	2 6 2
6	1 9 4	1 9 7	1 9 2	2 4	2 7 3	2 7 2	2 6 5
8	2 1 3	2 1 3	2 0 8	2 6	2 8 0	2 7 6	2 7 2
1 0	2 2 3	2 2 3	2 2 0	2 8	2 8 5	2 8 2	2 7 5
1 2	2 3 5	2 3 6	2 3 6	3 0	2 8 8	2 8 5	2 8 0
1 4	2 3 9	2 4 0	2 3 8	3 2	2 9 4	2 9 2	2 8 2
1 6	2 4 4	2 4 3	2 4 0	3 4	2 9 8	2 9 6	2 8 9
1 8	2 5 5	2 5 5	2 4 8	3 6	3 0 4	3 0 1	2 9 3

[0034]

[Table 4]

各群の平均飼料摂取量

[g/2匹/2日]

試験開始 から日目	対照食	0.5 M	5 M	試験開始 から日目	対照食	0.5 M	5 M
2	5 4	5 0	4 8	2 0	6 4	5 6	5 4
4	7 4	6 9	6 2	2 2	6 1	6 0	5 7
6	7 0	7 6	6 8	2 4	5 9	6 5	5 7
8	8 2	7 9	7 8	2 6	5 8	5 5	5 4
1 0	7 1	7 5	7 7	2 8	5 5	5 7	5 3
1 2	6 6	6 4	7 1	3 0	5 4	5 6	5 5
1 4	6 5	6 4	6 2	3 2	5 4	5 8	5 2
1 6	6 4	6 5	5 6	3 4	5 3	5 4	5 4
1 8	6 3	6 0	5 5	3 6	5 4	5 5	5 5

[0035]

Moreover, it collected blood from the subclavian vein in part to the heparin (15 units) content silicon coat micro spitz under anesthesia with the ether by the conditions of not abstaining from food, after [of experiment initiation] two weeks (about 1ml), and the blood serum was separated (2500rpm, 15 minutes, 4 degrees C). Moreover, blood collecting five weeks after was performed on the conditions of not abstaining from food, from experiment initiation, the ether performed it from the abdominal aorta under anesthesia, and the plasma of the same conditions as the above was obtained.

[0036]

Furthermore, measurement of the amount of growth hormones in the plasma obtained after [of experiment initiation] two weeks and the plasma obtained after five weeks used the rat growth hormone enzyme immunoassay (EIA) system (Amersham Life Science, England). And the amount of hormone in the plasma of two weeks after is shown in Table 5, and the amount of growth hormones in the plasma obtained after five weeks is shown in Table 6.

[0037]

[Table 5]

2 週間投与後の血漿中成長ホルモン濃度〔ng/ml〕		
対照食	0.5 M	5 M
92.3 ± 4.9	88.7 ± 8.0	83.6 ± 7.5

n=6、平均値±標準誤差

[0038]

[Table 6]

5 週間投与後の血漿中成長ホルモン濃度〔ng/ml〕		
対照食	0.5 M	5 M
103.2 ± 8.1	120.3 ± 7.8	120.6 ± 12.6

n=6、平均値±標準誤差

[0039]

As shown in Table 3 and 4 these results, the difference was not accepted, but adverse reaction was not accepted in whenever [growth-with Wistar-rats / which prescribed for the patient and bred the feed which added MAKa extractives powder /, and Wistar rats which prescribed for the patient and bred contrast feed], and (gain of body weight), the amount of feed intake at all, either, but growth with all three favorable groups was accepted in them.

[0040]

Moreover, although, as for the amount of growth hormones in the plasma after two weeks of administration, the inclination low a little was seen in the experimental group (group which prescribed MAKa for the patient) as shown in Table 5 and 6, the significant difference was not accepted in 3 between groups. Furthermore, as for the amount of growth hormones in the plasma obtained after five weeks of administration, as compared with the control group, the high value was accepted by the experimental group (0.5M, 5 M car group).

[0041]

Therefore, when it continued for five weeks to Wistar rats, they were supplied with the feed which added the water alcoholic extract dry matter of the dry matter of MAKa and it was bred to them, the rat was grown favorably and the growth hormone in the plasma in these Wistar rats increased. Consequently, it is safe for the body, there is no side effect, and by taking in the functional food which made the water alcoholic extract dry matter of the dry matter of MAKa contain as a principal component showed that the amount of growth hormones in blood rose.

[0042]

(Example 3 of an experiment)

Next, the effect affect the tenacity of the mouse by the ingestion of functional food which made the alcoholic extract extractives dry matter of MAKa of this invention contain is explained using the example of an experiment.

[0043]

First, the purpose which evaluates the operation over the whole living body function to the beginning considered the effect of the ingestion of MAKa to tenacity.

[0044]

The DDY mouse of 11 weeks old considered to have reached the adult mostly as a candidate for evaluation of tenacity was used. These DDY(s) mouse was used after arriving at a laboratory, and carrying out training

breeding for one week. Moreover, as tenacity, each DDY mouse was put in inside the cylinder with a diameter of 15cm, each [these] DDY mouse was run with constant speed, the cylinder was rotated, and time amount (distance) until it becomes impossible for each [these] DDY mouse to run with constant speed was evaluated as an index. In addition, the transit experiment was measured in the same time zone of the days.

[0045]

Moreover, after MAKKA used the dry matter of the root of MAKKA as impalpable powder, it was suspended so that it might become water in ml and 0.25mg /, and carried out direct single-dose administration of the 0.25ml by the sound into the stomach of each DDY mouse.

[0046]

Here, although it divided into the group of ten animals, having assumed whether a big difference would have any mice in a transit moving state and measurement was begun at first about selection of a mouse, when individual difference did not select by the preliminary transit test greatly in respect of lack of transit capacity or transit volition etc., it became clear by the simple group division that it could not use for an experiment.

[0047]

Then, the individual was first selected on the basis of being stabilized comparatively and always running in the mouse of the same weeks old, although a difference is in transit capacity. However, since it was difficult to make two mouse groups (a control group and intake group) which do not have a difference in normal transit capacity greatly, it was stabilized comparatively and the transit time in MAKKA administration order was always compared to the selected population it runs.

[0048]

Subsequently, as a result of investigating the effect which the capacity (ml) which prescribes MAKKA for the patient by the sound is changed about the effect the administration capacity of MAKKA affects intake of a diet, and it has on the intake of the diet of a mouse, when 0.5ml or more was prescribed for the patient, it became clear that the intake of a diet decreased and it was decided that administration capacity would be 0.1ml.

[0049]

Furthermore, change with the transit time before administration of MAKKA (distance) and the transit time during three days (distance) after prescribing 0.25ml of 0.1mg [/ml] MAKKA suspension for the patient was measured about seven mice (one-animal death) which the single-dose administration of MAKKA selected as mentioned above about the effect which it has on the endurance running of a mouse. consequently, it is shown in drawing 1 thru/or drawing 6 — as — the inside of six animals — four animals — the transit time — 24 hours after — setting — coming out — 50% or less 25% or more — it came out comparatively and increased. In addition, there was no change in two next animals. Moreover, although it was a reduction degree also after two days and three days, the somewhat high rate was shown. Furthermore, since it returned to the original rate after the 4th, it has been interpreted as maintaining the effectiveness of this amount of single-dose administration during the 3rd or less the 2nd day or more.

[0050]

From the result mentioned above, the effect which it has on the tenacity (endurance running) of the mouse by internal use of MAKKA was considered. In addition, in order to grasp an amount with exact intake, a medicine was prescribed for the patient using the approach of carrying out forcible administration by the sound. As a result of giving 0.025mg [per animal / /] MAKKA day by the administration capacity (0.1ml) which does not affect the diet intake of a mouse, among six animals, by four animals (66.6%), the increment in tenacity was accepted and the fall of tenacity was not shown in other two animals.

[0051]

Moreover, although it is as a result of four females and two males since one female died the middle among ten used mice (five males, five females), and three males did not become weak but it became measurement impossible for escape from a run state As shown in drawing 1 thru/or drawing 6 , among four animals, for the female, effectiveness was accepted by three animals (75%), and was accepted at a rate of one animal (50%) among two animals by the male.

[0052]

(Example 4 of an experiment)

Next, the effectiveness exerted on the tenacity of the mouse by the repetitive administration of the functional food which made the alcoholic extract extractives dry matter of MAKKA of this invention contain is explained using the example of an experiment.

[0053]

First, the same DDY mouse (two males and four females) as the above-mentioned example 3 of an experiment was prepared, the ingestion of the dry matter of the same MAKKA as the above-mentioned example 3 of an experiment was continued and carried out to each of each [these] mouse every count day of plurality, and repetitive administration was carried out.

[0054]

First, as shown in drawing 7 thru/or drawing 12 , the transit time for front [experiment initiation] three days was measured, and, specifically, the mean transit time for these three days was computed. In addition, the these-computed mean transit time was shown as a broken line in drawing 7 thru/or drawing 12 .

[0055]

Then, in each on the 11th and the 13th, specified quantity administration of the dry matter of MAKKA was carried out [the experiment opening day (the 0th day) and] on the 4th at each mouse, respectively. Here, in drawing 7 thru/or drawing 12 , the arrow head showed the day which prescribed the dry matter of MAKKA for the patient.

[0056]

Consequently, the mouse which mileage extended one day and two days after administration whenever it made the dry matter of MAKKA prescribe for the patient was observed. Extension of mileage has been checked on the 1st and the 2nd after prescribing the dry matter of MAKKA for the patient in four mice, the mouse (male 1) specifically shown in the inside of six mice shown in drawing 7 thru/or drawing 12 , and drawing 7 , the mouse (male 3) shown in drawing 9 , the mouse (female 5) shown in drawing 11 , and the mouse (female 6) shown in drawing 12 .

[0057]

Moreover, although the measurement of mileage of the mouse (female 4) shown in drawing 10 became impossible from the 6th day of experiment initiation on the way, it has checked stretch of mileage in administration of MAKKA in an experiment opening day.

[0058]

From the above thing, among six animals, whenever it prescribed MAKKA for the patient in five mice, improvement in tenacity has been checked by intake of MAKKA by having checked extension of mileage. Moreover, in each mouse other than the mouse (female 2) shown in drawing 8 , it has checked that there was a significant difference statistically by $P < 0.05$.

[0059]

(Example 1 of pharmaceutical preparation)

Citrus flavor 0.01g and 10g of potato starch were mixed to 1g of powder of the MAKKA extractives powder obtained in the example 1 of an experiment, and the tablet (food) was prepared with the conventional method.

[0060]

(Example 2 of pharmaceutical preparation)

Water was added to the thing of a presentation which added 10g of purified sucrose, and citrus flavor 0.05g to 0.5g of **** of the MAKKA extractives powder obtained in the example 1 of an experiment, it considered as 120ml of whole quantity, and the drink product (drinkable preparations) was prepared as plastics bottle stuffing.

[0061]

[Translation done.]

*** NOTICES ***

JPO and NCIP are not responsible for any damages caused by the use of this translation.

1. This document has been translated by computer. So the translation may not reflect the original precisely.
2. **** shows the word which can not be translated.
3. In the drawings, any words are not translated.

DESCRIPTION OF DRAWINGS

[Brief Description of the Drawings]

[Drawing 1] It is the graph which shows the example of an experiment which medicated the mouse (male 1) with the functional food containing MAKa of this invention.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 2] It is the graph which shows the example of an experiment which medicated the mouse (female 2) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 3] It is the graph which shows the example of an experiment which medicated the mouse (male 3) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 4] It is the graph which shows the example of an experiment which medicated the mouse (female 4) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 5] It is the graph which shows the example of an experiment which medicated the mouse (female 5) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 6] It is the graph which shows the example of an experiment which medicated the mouse (female 6) with the functional food containing MAKa same as the above.

- (a) The graph which shows change of the weight before and behind MAKa administration.
- (b) The graph which shows change of the food intake before and behind MAKa administration.
- (c) The graph which shows change of the movement tenacity before and behind MAKa administration.

[Drawing 7] It is the graph which shows the transit time of the mouse (male 1) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 8] It is the graph which shows the transit time of the mouse (female 2) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 9] It is the graph which shows the transit time of the mouse (male 3) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 10] It is the graph which shows the transit time of the mouse (female 4) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 11] It is the graph which shows the transit time of the mouse (female 5) which carried out repetitive administration of the functional food containing MAKa same as the above.

[Drawing 12] It is the graph which shows the transit time of the mouse (female 6) which carried out repetitive

administration of the functional food containing MAKAs same as the above.

[Translation done.]


* NOTICES *

JP0 and NCIP1 are not responsible for any damages caused by the use of this translation.


- 1.This document has been translated by computer. So the translation may not reflect the original precisely.
- 2.**** shows the word which can not be translated.
- 3.In the drawings, any words are not translated.

DRAWINGS


[Drawing 1]




[Drawing 2]



(a)




(b)




(c)


[Drawing 3]



(a)






(b)






(c)




[Drawing 4]




[Drawing 5]




[Drawing 6]




[Drawing 7]




[Drawing 8]




[Drawing 9]




[Drawing 10]



[Drawing 11]



[Drawing 12]



[Translation done.]

AN 2003:277167 CAPLUS
 DN 138:276809
 TI Studies on hygroscopic abilities of the water-extracted **Maca**
 AU Shimofuruya, Hiroshi; Suzuki, Ikukatsu; Funieda, Yoshihiko
 CS Suzuka Natl. Coll. Tech., Japan
 SO Kiyo - Suzuka Kogyo Koto Senmon Gakko (2003), 36, 131-134
 CODEN: SKSKDJ; ISSN: 0286-5483
 PB Suzuka Kogyo Koto Senmon Gakko
 DT Journal
 LA Japanese
 TI Studies on hygroscopic abilities of the water-extracted **Maca**
 AB Hygroscopic abilities of **Maca**, extd. from the powder of **Lepidium meyenii** Walp, were examd. by physicochem. techniques in comparison with those of urea, glycerol and D-glucitol used in cosmetics. When the relative humidity increased from 31.0% to 91.0%, the increased amts. of the moisture absorption capacity of urea and D-glucitol after 24 h standing were 85% and 66%, resp., whereas its amt. of the H2O-extd. **Maca** was 28%. The moisture absorption capacity of urea and D-glucitol were dependent on changes in the relative humidity, while the moisture absorption capacity of H2O-extd. **Maca** was not greatly influenced by changes in the relative humidity. Also, in the dry SiO2-gel desiccator, the H2O-extd. **Maca** showed the best moisture retention capacity among the samples tested. Probably the H2O-extd. **Maca** was a desirable hygroscopic material because it exhibited relative high hygroscopic ability under conditions with various humidity and high moisture retention capacity even in the dry SiO2-gel desiccator.
 ST humidity hygroscopic ability water extd **Maca**
 IT Cosmetics
 Hygroscopic substances
 Lepidium peruvianum
 (hygroscopic abilities of water-extd. **Maca**)
 IT Humidity